

GARS Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5648

Product Information

Application	WB
Primary Accession	P41250
Other Accession	Q5RBL1
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	83166
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	2617
Antigen Region	706-739
Other Names	Glycine--tRNA ligase, Diadenosine tetraphosphate synthetase, AP-4-A synthetase, Glycyl-tRNA synthetase, GlyRS, GARS
Dilution	WB~~1:4000
Target/Specificity	This GARS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 706-739 amino acids from the C-terminal region of human GARS.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GARS Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GARS1 (HGNC:4162)
Synonyms	GARS

Function	Catalyzes the ATP-dependent ligation of glycine to the 3'-end of its cognate tRNA, via the formation of an aminoacyl-adenylate intermediate (Gly-AMP) (PubMed: 17544401 , PubMed: 24898252 , PubMed: 28675565). Also produces diadenosine tetraphosphate (Ap4A), a universal pleiotropic signaling molecule needed for cell regulation pathways, by direct condensation of 2 ATPs. Thereby, may play a special role in Ap4A homeostasis (PubMed: 19710017).
Cellular Location	Cytoplasm. Cell projection, axon. Secreted {ECO:0000250 UniProtKB:Q9CZD3}. Secreted, extracellular exosome {ECO:0000250 UniProtKB:Q9CZD3}. Note=In transfected COS7 cells, not detected in mitochondria, nor in Golgi apparatus (PubMed:17035524) Secreted by motor neuron, possibly through the exosome pathway (By similarity). {ECO:0000250 UniProtKB:Q9CZD3, ECO:0000269 PubMed:17035524} [Isoform 2]: Cytoplasm. Cell projection, axon
Tissue Location	Widely expressed, including in brain and spinal cord. [Isoform 1]: Expressed in brain, spinal cord, muscle, heart, spleen and liver.


Background

GARS is a glycyl-tRNA synthetase, one of the aminoacyl-tRNA synthetases that charge tRNAs with their cognate amino acids. This protein is an (alpha)₂ dimer which belongs to the class II family of tRNA synthetases. The protein has been shown to be a target of autoantibodies in the human autoimmune diseases, polymyositis or dermatomyositis.

References

Shiba K., Schimmel P.J. Biol. Chem. 269:30049-30055(1994) Antonellis A., Ellsworth R.E.Am. J. Hum. Genet. 72:1293-1299(2003)

Images



All lanes : Anti-GARS Antibody (C-term) at 1:2000 dilution
 Lane 1: NCI-H460 whole cell lysate Lane 2: HT-1080 whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: C6 whole cell lysate Lane 5: Ramos whole cell lysate
 Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 83 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.