

ACO1 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AW5676

Product Information

Application	WB
Primary Accession	P21399
Other Accession	Q01059 , Q63270
Reactivity	Human, Mouse, Rat
Predicted	Chicken
Host	Rabbit
Clonality	Polyclonal
Calculated MW	98399
Isotype	Rabbit IgG
Antigen Source	HUMAN

Additional Information

Gene ID	48
Antigen Region	124-155
Other Names	Cytoplasmic aconitate hydratase, Aconitase, 4.2.1.3, Citrate hydro-lyase, Ferritin repressor protein, Iron regulatory protein 1, IRP1, Iron-responsive element-binding protein 1, IRE-BP 1, ACO1, IREB1
Dilution	WB~~1:1000
Target/Specificity	This ACO1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 124-155 amino acids from human ACO1.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ACO1 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ACO1
-------------	------

Synonyms

IREB1

Function

Bifunctional iron sensor that switches between 2 activities depending on iron availability (PubMed:[1281544](#), PubMed:[1946430](#), PubMed:[8041788](#)). Iron deprivation, promotes its mRNA binding activity through which it regulates the expression of genes involved in iron uptake, sequestration and utilization (PubMed:[1281544](#), PubMed:[1946430](#), PubMed:[23891004](#), PubMed:[8041788](#)). Binds to iron-responsive elements (IRES) in the untranslated region of target mRNAs preventing for instance the translation of ferritin and aminolevulinic acid synthase and stabilizing the transferrin receptor mRNA (PubMed:[1281544](#), PubMed:[1946430](#), PubMed:[23891004](#), PubMed:[8041788](#)).

Cellular Location

Cytoplasm, cytosol.


Background

Iron sensor. Binds a 4Fe-4S cluster and functions as aconitase when cellular iron levels are high. Functions as mRNA binding protein that regulates uptake, sequestration and utilization of iron when cellular iron levels are low. Binds to iron-responsive elements (IRES) in target mRNA species when iron levels are low. Binding of a 4Fe-4S cluster precludes RNA binding.

References

- Hirling H.,et al.Nucleic Acids Res. 20:33-39(1992).
Humphray S.J.,et al.Nature 429:369-374(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Rouault T.A.,et al.Proc. Natl. Acad. Sci. U.S.A. 87:7958-7962(1990).
Hentze M.W.,et al.Nucleic Acids Res. 19:1739-1740(1991).

Images



All lanes : Anti-ACO1 Antibody (N-Term) at 1:1000 dilution
Lane 1: mouse kidney lysate Lane 2: rat kidney lysate
Lane 3: U-2OS whole cell lysate Lane 4: human kidney lysate Lane 5: mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 98 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.