

DANRE heg Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # Azb10023b

Product Information

Application WB, E **Primary Accession Q6R8|2** Zebrafish Reactivity Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Calculated MW** 105529 **Antigen Region** 691-717

Additional Information

Other Names Protein HEG, Heart of glass, heg

Target/Specificity This DANRE heg antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 691-717 amino acids from the

C-terminal region of DANRE heg.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions DANRE heg Antibody (C-term) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name heg

Function Receptor component of the CCM signaling pathway which is a crucial

regulator of heart and vessel formation and integrity. May act through the

stabilization of endothelial cell junctions.

Cellular Location [Isoform 1]: Cell membrane; Single- pass type I membrane protein. Cell

junction [Isoform 3]: Secreted.


Background

Receptor component of the CCM signaling pathway which is a crucial regulator of heart and vessel formation and integrity. May act through the stabilization of endothelial cell junctions.

References

Mably J.D., et al. Curr. Biol. 13:2138-2147(2003). Kleaveland B., et al. Nat. Med. 15:169-176(2009).

Images



All lanes: Anti-(DANRE) heg Antibody (C-term) at 1:1000 dilution Lane 1: Zebrafish muscle lysate Lane 2: Zebrafish lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 106 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.