

# DANRE gnb1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # Azb10033a

#### **Product Information**

**Application** WB, E **Primary Accession Q6PH57** Reactivity Zebrafish Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Calculated MW** 37286 **Antigen Region** 1-30

#### **Additional Information**

**Gene ID** 322104

Other Names Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1, Transducin

beta chain 1, gnb1

Target/Specificity This DANRE gnb1 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 1-30 amino acids from the N-terminal

region of DANRE gnb1.

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** DANRE gnb1 Antibody (N-term) is for research use only and not for use in

diagnostic or therapeutic procedures.

### **Protein Information**

Name gnb1

**Function** Guanine nucleotide-binding proteins (G proteins) are involved as a


modulator or transducer in various transmembrane signaling systems. The beta and gamma chains are required for the GTPase activity, for replacement

of GDP by GTP, and for G protein-effector interaction (By similarity).


## **Background**

Guanine nucleotide-binding proteins (G proteins) are involved as a modulator or transducer in various transmembrane signaling systems. The beta and gamma chains are required for the GTPase activity, for replacement of GDP by GTP, and for G protein-effector interaction (By similarity).

## **Images**



All lanes: Anti-(DANRE) gnb1 Antibody (center) at 1:1000 dilution Lane 1: Zebrafish lysate Lane 2: ZF4 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 38 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



DANRE gnb1 Antibody (N-term) (Cat. #Azb10033a) western blot analysis in zebra fish brain tissue lysates (35ug/lane). This demonstrates the DANRE gnb1 antibody detected the DANRE gnb1 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.