

# (DANRE) znf148 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # Azb18713c

#### **Product Information**

**Application** WB, E **Primary Accession** A0MS83 Reactivity Zebrafish Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB46462 **Calculated MW** 90507

#### **Additional Information**

**Gene ID** 560300

Other Names Zinc finger protein 148, Transcription factor ZBP-89, Zinc finger DNA-binding

protein 89, znf148, zbp89

**Target/Specificity** This (DANRE) znf148 antibody is generated from a rabbit immunized with a

KLH conjugated synthetic peptide between 287-320 amino acids of DANRE

znf148.

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** (DANRE) znf148 Antibody (Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

## **Protein Information**

Name znf148

Synonyms zbp89

**Function** Involved in transcriptional regulation. Represses the transcription of a

number of genes. Required for primitive and definitive hematopoiesis during

embryonic development.


# **Background**

Involved in transcriptional regulation. Represses the transcription of a number of genes. Required for primitive and definitive hematopoiesis during embryonic development.

## References

Li X.,et al.Development 133:3641-3650(2006). Howe K.,et al.Nature 496:498-503(2013).

## **Images**



Western blot analysis of lysates from zebra fish muscle, Zebrafish tissue lysate (from left to right), using (DANRE) znf148 Antibody (Center)(Cat. #Azb18713c). Azb18713c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.