

(DANRE) opn1sw2 Antibody (C-Term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # Azb21565b

Product Information

Application	WB, E
Primary Accession	Q9W6A8
Reactivity	Zebrafish
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Clone Names	RB52986
Calculated MW	39484

Additional Information

Gene ID	30435
Other Names	Opsin-1, short-wave-sensitive 2, Blue cone photoreceptor pigment, Blue-sensitive opsin, Opsin SWS-2, opn1sw2, bluops, opn1sw1, sws2
Target/Specificity	This DANRE opn1sw2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 242-276 amino acids of DANRE opn1sw2.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	(DANRE) opn1sw2 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	opn1sw2
Synonyms	bluops, opn1sw1, sws2
Function	Visual pigments are the light-absorbing molecules that mediate vision. They consist of an apoprotein, opsin, covalently linked to cis-retinal.

Cellular Location Membrane; Multi-pass membrane protein.

Tissue Location Retinal long single cone outer segments.

Background

Visual pigments are the light-absorbing molecules that mediate vision. They consist of an apoprotein, opsin, covalently linked to cis-retinal.


References

Vihtelic T.S., et al. *Vis. Neurosci.* 16:571-585(1999).

Chinen A., et al. *Genetics* 163:663-675(2003).

Howe K., et al. *Nature* 496:498-503(2013).

Images



Anti-opn1sw2 Antibody (C-Term) at 1:2000 dilution + ZF4 whole cell lysates. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 39 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

Citations

- [Deficiency of copper responsive gene stm4 induces retinal developmental defects.](#)
- [Silver nanoparticles affect lens rather than retina development in zebrafish embryos.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.