10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



GM-CSF

Catalog # PVGS1022

Product Information

Primary Accession P04141
Species Human

Sequence Ala18-Glu144, expressed with an N-terminal Met

Purity > 95% as analyzed by SDS-PAGE

Endotoxin Level

Expression System P. pastoris

Formulation Lyophilized after extensive dialysis against 10 mM PB, pH 7.0.

Reconstitution It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

ddH₂O up to 100 □g/ml.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at lower than

-70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw

cycles.

Additional Information

Gene ID 1437

Other Names Granulocyte-macrophage colony-stimulating factor, GM-CSF,

Colony-stimulating factor, CSF, Molgramostin, Sargramostim, CSF2, GMCSF

Target Background Granulocyte Macrophage-Colony Stimulating Factor (GM-CSF) was initially

characterized as a growth factor that can support the in vitro colony

formation of granulocyte-macrophage progenitors. Granulocyte

Macrophage-Colony Stimulating Factor (GM-CSF) is produced by a number of different cell types, including activated T cells, B cells, macrophages, mast cells, endothelial cells, and fibroblasts, in response to cytokine of immune and

inflammatory stimuli. Besides granulocyte-macrophage progenitors, Granulocyte Macrophage-Colony Stimulating Factor (GM-CSF) is a growth factor for erythroid, megakaryocyte, and eosinophil progenitors. On mature

hematopoietic, monocytes/macrophages and eosinophils. Human

Granulocyte Macrophage-Colony Stimulating Factor (GM-CSF) can induce human endothelial cells to migrate and proliferate. Additionally, Granulocyte

Macrophage-Colony Stimulating Factor (GM-CSF) can stimulate the

proliferation of a number of tumor cell lines, including osteogenic sarcoma,

carcinoma, and adenocarcinoma cell lines.

Protein Information

Name CSF2

Synonyms GMCSF

Function Cytokine that stimulates the growth and differentiation of hematopoietic

precursor cells from various lineages, including granulocytes, macrophages,

eosinophils and erythrocytes.

Cellular Location Secreted

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.