

FGF-basic

Catalog # PVGS1038

Product Information

Species Rat

Sequence MPALPEDGGG AFPPGHFKDP KRLYCKNGGF FLRIHPDGRV DGVREKSDPH

VKLQLQAEER GVVSIKGVCA NRYLAMKEDG RLLASKCVTE ECFFFERLES NNYNTYRSRK YSSWYVALKR TGQYKLGSKT GPGQKAILFL PMSAKS

Purity > 98 % by SDS-PAGE and HPLC analyses.

Endotoxin Level Less than 1 EU/ Leg of rRtbFGF as determined by LAL method.

Formulation Lyophilized from a 0.2 \(\text{Im filtered solution in PBS, pH 7.4.} \)

Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring

the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock

solutions should be apportioned into working aliquots and stored at \leq -20 °C.

Further dilutions should be made in appropriate buffered solutions.

Additional Information

Target Background

FGF-basic is one of 23 known members of the FGF family. Proteins of this family play a central role during prenatal development and postnatal growth and regeneration of a variety of tissues, by promoting cellular proliferation and differentiation. FGF-basic is a non-glycosylated heparin binding growth factor that is expressed in the brain, pituitary, kidney, retina, bone, testis, adrenal gland liver, monocytes, epithelial cells and endothelial cells. FGF-basic signals through FGFR 1b, 1c, 2c, 3c and 4.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.