10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

EMAP-II, Human

Catalog # PVGS1048

Product Information

Species	Human
Sequence	SKPIDVSRLD LRIGCIITAR KHPDADSLYV EEVDVGEIAP RTVVSGLVNH VPLEQMQNRM VILLCNLKPA KMRGVLSQAM VMCASSPEKI EILAPPNGSV PGDRITFDAF PGEPDKELNP KKKIWEQIQP DLHTNDECVA TYKGVPFEVK GKGVCRAQTM SNSGIK
Purity	> 98 % by SDS-PAGE and HPLC analyses.
Endotoxin Level	Less than 1 EU/ μg of rHuEMAP-II as determined by LAL method.
Formulation **Reconstitution**	Lyophilized from a 0.2 μm filtered concentrated solution in PBS, pH 7.4. We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Additional Information

Target Background	EMAP-II is a tumor derived cytokine that exerts a wide range of activities on endothelial cells, monocytes and neutrophils. EMAP-II inhibits endothelial cell proliferation, vasculogenesis, neovessel formation, and can induce apoptosis. It is also chemotactic towards neutrophils and monocytes and induces myeloperoxidase activity from neutrophils. Of clinical importance, EMAP-II inhibits angiogenesis of vascular beds and suppresses the growth of primary and secondary tumors without affecting normal tissues. Mature EMAP-II is an 18.3 kDa protein, which is synthesized as the C-terminal portion of a biologically inactive precursor protein containing a propeptide of 146 amino acid residues.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.