

GM-CSF

Catalog # PVGS1052

Product Information

Primary Accession 09GL44

Species Rhesus Macaque

Sequence Ala18-Glu144

Purity > 98% as analyzed by SDS-PAGE

> 98% as analyzed by HPLC

Endotoxin Level

Biological Activity Fully biologically active when compared to standard. The ED₅₀ as determined

by a cell proliferation assay using human TF-1 cells is less than 0.1 ng/ml,

corresponding to a specific activity of $> 1.0 \times 10^7$ IU/mg.

Expression System E. coli

Theoretical Molecular Weight 14.4 kDa

FormulationLyophilized from a 0.2 Im filtered solution in PBS, pH 7.4. **Reconstitution**It is recommended that this vial be briefly centrifuged prio

on It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in

sterile distilled water or aqueous buffer containing 0.1% BSA to a

concentration of 0.1-1.0 mg/ml.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at -70°C or

-20°C. Upon reconstitution, the product should be stable for up to 1 week at

4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Additional Information

Target Background Granulocyte Macrophage-Colony Stimulating Factor (GM-CSF) was initially

characterized as a growth factor that can support the in vitro colony formation of granulocyte-macrophage progenitors. Granulocyte

Macrophage-Colony Stimulating Factor (GM-CSF) is produced by a number of different cell types, including activated T cells, B cells, macrophages, mast cells, endothelial cells, and fibroblasts, in response to cytokine of immune and

inflammatory stimuli. Besides granulocyte-macrophage progenitors, Granulocyte Macrophage-Colony Stimulating Factor (GM-CSF) is a growth factor for erythroid, megakaryocyte, and eosinophil progenitors. On mature hematopoietic, monocytes/macrophages and eosinophils. Additionally, Granulocyte Macrophage-Colony Stimulating Factor (GM-CSF) can stimulate the proliferation of a number of tumor cell lines, including osteogenic

sarcoma, carcinoma, and adenocarcinoma cell lines.

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.