10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

BCMA

Catalog # PVGS1063

Product Information

Primary Accession	Q02223
Species	Human
Sequence	Ala5-Ala54
Purity	> 98% as analyzed by SDS-PAGE > 98% as analyzed by HPLC
Endotoxin Level	
Biological Activity	Fully biologically active when compared to standard. The ED_{50} as determined by its ability to inhibit APRIL-mediated proliferation of anti-IgM stimulated murine B cells is no less than 40.0 ng/ml, corresponding to a specific activity of > 2.5×10^4 IU/mg in the presence of 100.0 ng/ml human APRIL.
Expression System	E. coli
Theoretical Molecular Weight	5.4 kDa
Formulation	Lyophilized from a 0.2 □m filtered solution in 30% acetonitrile, 0.1% TFA.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	608
Other Names	Tumor necrosis factor receptor superfamily member 17, B-cell maturation protein, CD269, TNFRSF17, BCM, BCMA
Target Background	BCMA, a member of the TNF receptor superfamily, binds to BAFF and APRIL. BCMA is expressed on mature B-cells and other B-cell lines and plays an important role in B cell development, function and regulation. BCMA also has the capability to activate NF-kappaB and JNK. The human BCMA gene codes for a 184 amino acid type I transmembrane protein, which contains a 54 amino acid extracellular domain, a 23 amino acid transmembrane domain, and a 107 amino acid extracellular domain.

Protein Information

Name	TNFRSF17
Synonyms	BCM, BCMA
Function	Receptor for TNFSF13B/BLyS/BAFF and TNFSF13/APRIL. Promotes B-cell survival and plays a role in the regulation of humoral immunity. Activates NF-kappa-B and JNK.
Cellular Location	Cell membrane; Single-pass type III membrane protein. Endomembrane system; Single-pass type III membrane protein Note=Perinuclear Golgi-like structures
Tissue Location	Expressed in mature B-cells, but not in T-cells or monocytes

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.