

FGF-19

Catalog # PVGS1072

Product Information

Primary Accession Species	O95750 Human
Sequence	Arg23-Lys216, expressed with an N-terminal Met
Purity	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
Endotoxin Level	
Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using murine Balb/c 3T3 cells is less than 150.0 ng/ml, corresponding to a specific activity of $> 6.7 \times 10^3$ IU/mg.
Expression System	E. coli
Theoretical Molecular Weight	21.8 kDa
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS, pH 7.4.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	9965
Other Names	Fibroblast growth factor 19, FGF-19, FGF19
Target Background	Fibroblast growth factor 19 (FGF19) belongs to the large FGF family which has at least 23 members. All FGF family members are heparin binding growth factors with a core 120 amino acid (aa) FGF domain that allows for a common tertiary structure. FGFs are expressed during embryonic development and in restricted adult tissues. Four distinct but related classes of FGF receptors, FGF R1, 2, 3, and 4, exist. Unlike most FGFs which bind to and activate more than one FGF receptor, FGF19 is a specific ligand for FGF R4.

Protein Information

Name	FGF19
Function	Involved in the suppression of bile acid biosynthesis through down-regulation of CYP7A1 expression, following positive regulation of the JNK and ERK1/2 cascades. Stimulates glucose uptake in adipocytes. Activity requires the presence of KLB and FGFR4.
Cellular Location	Secreted.
Tissue Location	Expressed in fetal brain, cartilage, retina, and adult gall bladder.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.