10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

BD-3

Catalog # PVGS1074

Product Information

Primary Accession	P81534
Species	Human
Sequence	Gly23-Lys67
Purity	> 98% as analyzed by SDS-PAGE > 98% as analyzed by HPLC
Endotoxin Level	
Biological Activity	Fully biologically active when compared to standard. The ED_{50} as determined by anti-microbial activity against E.coli is less than 30.0 g/ml, corresponding to a specific activity of > 33.3 IU/mg.
Expression System	E. coli
Theoretical Molecular Weight	5.2 kDa
Formulation	Lyophilized from a 0.2 m filtered solution in 20 mM PBS, pH 7.4, 130 mM NaCl.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	414325;55894
Other Names	Beta-defensin 103, Beta-defensin 3, BD-3, DEFB-3, HBD3, hBD-3, Defensin, beta 103, Defensin-like protein, DEFB103A, BD3, DEFB103, DEFB3
Target Background	Defensins (alpha and beta) are cationic peptides with a broad spectrum of antimicrobial activity that comprise an important arm of the innate immune system. The α-defensins are distinguished from the β-defensins by the pairing of their three disulfide bonds. To date, four human β-defensins have been identified; BD-1, BD-2, BD-3 and BD-4. β-defensins are expressed on some leukocytes and at epithelial surfaces. In addition to their direct antimicrobial activities, they are chemoattractant towards immature dendritic cells and memory T cells. The β-defensin proteins are expressed as the C-terminal portion of precursors and are released by proteolytic cleavage of a signal

sequence and, in the case of BD-1 (36 a.a.), a propeptide region. β-defensins contain a six-cysteine motif that forms three intra-molecular disulfide bonds. β-Defensins are 3-5 kDa peptides ranging in size from 33-47 amino acid residues.

Protein Information

Name	DEFB103A
Synonyms	BD3, DEFB103, DEFB3
Function	Exhibits antimicrobial activity against Gram-positive bacteria S.aureus and S.pyogenes, Gram-negative bacteria P.aeruginosa and E.coli and the yeast C.albicans. Kills multiresistant S.aureus and vancomycin-resistant E.faecium. No significant hemolytic activity was observed.
Cellular Location	Secreted.
Tissue Location	Highly expressed in skin and tonsils, and to a lesser extent in trachea, uterus, kidney, thymus, adenoid, pharynx and tongue. Low expression in salivary gland, bone marrow, colon, stomach, polyp and larynx. No expression in small intestine

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.