

Lymphotactin/XCL1

Catalog # PVGS1092

Product Information

Primary Accession P47992
Species Human

Sequence Gly23-Gly114

Purity > 98% as analyzed by SDS-PAGE

> 98% as analyzed by HPLC

Endotoxin Level

Biological Activity Fully biologically active when compared to standard. The biological activity

determined by a chemotaxis bioassay using human T-lymphocytes is in a

concentration of 10.0-100.0 ng/ml.

Expression System E. coli

Theoretical Molecular Weight 10.2 kDa

Formulation Lyophilized from a 0.2 Im filtered solution in 20 mM PB, pH 7.4, 150 mM

NaCl.

Reconstitution It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1 % BSA to a

concentration of 0.1-1.0 mg/ml.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at -70°C or

-20°C. Upon reconstitution, the product should be stable for up to 1 week at

4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID 6375

Other Names Lymphotactin, ATAC, C motif chemokine 1, Cytokine SCM-1, Lymphotaxin,

SCM-1-alpha, Small-inducible cytokine C1, XC chemokine ligand 1, XCL1, LTN,

SCYC1

Target Background Lymphotactin is the only known member of the C-chemokine family and

signals through the receptor XCR1, formally known as GPR5. The spleen shows the highest level of lymphotactin compared to peripheral leukocytes,

lung, colon and small intestine. Lymphotactin is chemotactic towards

lymphocytes but not towards monocytes or neutrophils.

Protein Information

Name XCL1

Synonyms LTN, SCYC1

Function Chemotactic activity for lymphocytes but not for monocytes or neutrophils.

In thymus, mediates medullary accumulation of thymic dendritic cells and contributes to regulatoy T cell development, playing a role in self-tolerance

establishment.

Cellular Location Secreted.

Tissue Location Highest level in spleen, lower in peripheral leukocytes and very low levels in

lung, colon and small intestine

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.