

# Exodus-2/CCL21

Catalog # PVGS1100

## Product Information

---

<b>Primary Accession Species</b>	<a href="#">P86792</a> Mouse
<b>Sequence</b>	Ser24-Gly133
<b>Purity</b>	> 97% as analyzed by SDS-PAGE > 97% as analyzed by HPLC
<b>Endotoxin Level Biological Activity</b>	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using murine T-lymphocytes is in a concentration range of 10.0-100.0 ng/ml.
<b>Expression System</b>	E. coli
<b>Theoretical Molecular Weight</b>	12.1 kDa
<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution in 20 mM PB, pH 7.4, 150 mM NaCl.
<b>Reconstitution</b>	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
<b>Storage &amp; Stability</b>	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

## Additional Information

---

<b>Gene ID</b>	100042493;100504239;100504346;100862177
<b>Other Names</b>	C-C motif chemokine 21b, 6Ckine, Beta-chemokine exodus-2, Small-inducible cytokine A21b, Thymus-derived chemotactic agent 4, TCA4, Ccl21b, Scya21, Scya21b
<b>Target Background</b>	Exodus-2/CCL21 is a novel CC chemokine discovered independently by three groups from the EST database, and shows 21-33% identity to other CC chemokines. Exodus-2 contains the four conserved cysteines characteristic of $\beta$ chemokines plus two additional cysteines in its unusually long carboxyl terminal domain. It is expressed in lymph nodes of certain endothelial cells, and in the spleen and appendix. Exodus-2 chemoattracts T and B lymphocytes and inhibits hematopoiesis.

## Protein Information

---

<b>Name</b>	Ccl21b
<b>Synonyms</b>	Scya21, Scya21b
<b>Function</b>	Inhibits hemopoiesis and stimulates chemotaxis. Chemotactic in vitro for thymocytes and activated T-cells, but not for B-cells, macrophages, or neutrophils. Potent mesangial cell chemoattractant. Shows preferential activity towards naive T-cells. May play a role in mediating homing of lymphocytes to secondary lymphoid organs.
<b>Cellular Location</b>	Secreted.
<b>Tissue Location</b>	Expressed strongly in lung, spleen, thymus, peripheral and mesenteric lymph nodes. Also expressed in the testis, kidney, liver, and heart

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.