

Eotaxin-2/CCL24

Catalog # PVGS1124

Product Information

Primary Accession Species	O00175 Human
Sequence	Val27-Ala104
Purity	> 97% as analyzed by SDS-PAGE > 97% as analyzed by HPLC
Endotoxin Level Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human peripheral blood eosinophils is in a concentration of 50.0-100.0 ng/ml.
Expression System	E. coli
Theoretical Molecular Weight	8.8 kDa
Formulation	Lyophilized from a 0.2 μ m filtered solution in 20 mM PB, pH 7.4, 150 mM NaCl.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	6369
Other Names	C-C motif chemokine 24, CK-beta-6, Eosinophil chemotactic protein 2 {ECO:0000303 Ref.6}, Eotaxin-2 {ECO:0000303 Ref.6}, Myeloid progenitor inhibitory factor 2, MPIF-2, Small-inducible cytokine A24, CCL24 (HGNC:10623)
Target Background	Eotaxin-2/CCL24, also named MPIF-2 and Ck β 6, is a novel CC chemokine recently identified. It is produced by activated monocytes and T lymphocytes. Eotaxin-2 selectively chemoattracts cells expressing CCR3 including eosinophils, basophils, Th2 T cells, mast cells, and certain subsets of dendritic cells. Additionally, Eotaxin-2 inhibits the proliferation of multipotential hematopoietic progenitor cells. The mature protein, which also includes a C-terminal truncation, contains 78 amino acid residues (92 a.a. residues for the mouse homolog, without C-terminal truncation).

Protein Information

Name	CCL24 (HGNC:10623)
Function	Chemotactic for resting T-lymphocytes, and eosinophils (PubMed: 9104803 , PubMed: 9365122). Has lower chemotactic activity for neutrophils but none for monocytes and activated lymphocytes (PubMed: 9104803 , PubMed: 9365122). Is a strong suppressor of colony formation by a multipotential hematopoietic progenitor cell line (PubMed: 9104803 , PubMed: 9365122). Binds to CCR3 (PubMed: 9104803 , PubMed: 9365122).
Cellular Location	Secreted
Tissue Location	Activated monocytes and activated T lymphocytes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.