

IL-1 α

Catalog # PVGS1152

Product Information

Primary Accession Species	Q3U0Y6 Mouse
Sequence	Ser115-Ser270
Purity	> 97% as analyzed by SDS-PAGE > 97% as analyzed by HPLC
Endotoxin Level Biological Activity	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using murine D10S cells is less than 2.0 pg/ml, corresponding to a specific activity of > 5.0 × 10 ⁸ IU/mg.
Expression System	E. coli
Theoretical Molecular Weight	17.9 kDa
Formulation Reconstitution	Lyophilized from a 0.2 μ m filtered solution in PBS, pH 7.4. It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Additional Information

Target Background	Interleukin-1 alpha (IL-1 α) is a non-secreted proinflammatory cytokine produced in a variety of cells including monocytes, tissue macrophages, keratinocytes and other epithelial cells. Both IL-1 α and IL-1 β binds to the same receptor and has similar if not identical biological properties. These cytokines have a broad range of activities including, stimulation of thymocyte proliferation, by inducing IL-2 release, B-cell maturation and proliferation, mitogenic FGF-like activity and the ability to stimulate the release of prostaglandin and collagenase from synovial cells. However, whereas IL-1 β is a secreted cytokine, IL-1 α is predominantly a cell-associated cytokine.
--------------------------	---

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.