

# HB-EGF

Catalog # PVGS1164

## Product Information

---

<b>Primary Accession Species</b>	<a href="#">Q06186</a> Mouse
<b>Sequence</b>	Asp63-Leu148
<b>Purity</b>	> 97% as analyzed by SDS-PAGE > 97% as analyzed by HPLC
<b>Endotoxin Level</b>	
<b>Biological Activity</b>	Fully biologically active when compared to standard. The ED <sub>50</sub> as determined by a cell proliferation assay using murine Balb/c 3T3 cells is less than 1.0 ng/ml, corresponding to a specific activity of > 1.0 × 10 <sup>6</sup> IU/mg.
<b>Expression System</b>	E. coli
<b>Theoretical Molecular Weight</b>	9.8 kDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution in 10 mM PB, 500 mM NaCl, pH7.4. It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
<b>Reconstitution</b>	
<b>Storage &amp; Stability</b>	Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

## Additional Information

---

<b>Gene ID</b>	15200
<b>Other Names</b>	Proheparin-binding EGF-like growth factor, Heparin-binding EGF-like growth factor, HB-EGF, HBEGF, Hbegf, Dtr, Hegfl
<b>Target Background</b>	Heparin-binding EGF-like growth factor (HB-EGF) is a member of the EGF family of proteins. HB-EGF-like growth factor is synthesized as a membrane-anchored mitogenic and chemotactic glycoprotein. An epidermal growth factor produced by monocytes and macrophages, due to an affinity for heparin is termed HB-EGF. It has been shown to play a role in wound healing, cardiac hypertrophy and heart development and function. The transmembrane form of HB-EGF is the unique receptor for diphtheria toxin and functions in juxtacrine signaling in cells. Both forms of HB-EGF participate in normal physiological processes and in pathological processes including tumor progression and metastasis, organ hyperplasia, and atherosclerotic disease.

HB-EGF can bind two locations on cell surfaces, heparan sulfate proteoglycans and EGF-receptor effecting cell to cell interactions.

## Protein Information

---

<b>Name</b>	Hbegf
<b>Synonyms</b>	Dtr, Hegfl
<b>Function</b>	Growth factor that mediates its effects via EGFR, ERBB2 and ERBB4. Required for normal cardiac valve formation and normal heart function. Promotes smooth muscle cell proliferation. May be involved in macrophage-mediated cellular proliferation. It is mitogenic for fibroblasts, but not endothelial cells. It is able to bind EGF receptor/EGFR with higher affinity than EGF itself and is a far more potent mitogen for smooth muscle cells than EGF. Also acts as a diphtheria toxin receptor.
<b>Cellular Location</b>	[Heparin-binding EGF-like growth factor]: Secreted, extracellular space. Note=Mature HB-EGF is released into the extracellular space and probably binds to a receptor
<b>Tissue Location</b>	Most abundant in kidney, skeletal muscle, lung, spleen, brain and heart

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.