

# TGIF; IL-10A

Catalog # PVGS1197

## Product Information

---

<b>Primary Accession</b>	<a href="#">P29456</a>
<b>Species</b>	Rat
<b>Sequence</b>	MSKGHSIKGD NNCTHFPVSQ THMLRELRAA FSQVKTFQK KDQLDNIVLT DSLLQDFKGY LGCQALSEMI KFYLVEMPQ AENHGPEIKE HLNSLGEKLG TLWIQLRRCH RFLPCENKSK AVEQVKNDFN KLQDKGVYKA MNEFDIFINC IEAYVTLKMK N
<b>Purity</b>	> 95% by SDS-PAGE and HPLC analyses.
<b>Endotoxin Level</b>	
<b>Formulation</b>	Lyophilized after extensive dialysis against PBS.
<b>Reconstitution</b>	Reconstituted in ddH <sub>2</sub> O at 100 µg/ml.

## Additional Information

---

<b>Gene ID</b>	25325
<b>Other Names</b>	Interleukin-10, IL-10, Cytokine synthesis inhibitory factor, CSIF, Il10, IL-10
<b>Target Background</b>	<p>Interleukin-10 (IL-10), also known as cytokine synthesis inhibitory factor (CSIF), is an anti-inflammatory cytokine produced by a variety of cell lines including T-cells, macrophages and mast cells. IL-10 is classified as a class-2 cytokine, a set of cytokines including IL-19, IL-20, IL-22, IL-24, and IL-26. IL-10 can inhibit the synthesis of pro-inflammatory cytokines such as IFN-gamma, IL-2, IL-3, TNF and GM-CSF. It also stimulates Th2 responses, but suppresses the antigen-presentation capacity of antigen presenting cells.</p> <p>Recombinant rat Interleukin-10 (IL-10) produced in E. coli is a single non-glycosylated polypeptide chain containing 161 amino acids. A fully biologically active molecule, recombinant rat Interleukin-10 (IL-10) has a molecular mass of 18.7kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at .</p>

## Protein Information

---

<b>Name</b>	Il10
<b>Synonyms</b>	IL-10
<b>Function</b>	Major immune regulatory cytokine that acts on many cells of the immune system where it has profound anti-inflammatory functions, limiting excessive

tissue disruption caused by inflammation. Mechanistically, IL10 binds to its heterotetrameric receptor comprising IL10RA and IL10RB leading to JAK1 and STAT2-mediated phosphorylation of STAT3. In turn, STAT3 translocates to the nucleus where it drives expression of anti-inflammatory mediators. Targets antigen-presenting cells (APCs) such as macrophages and monocytes and inhibits their release of pro-inflammatory cytokines including granulocyte-macrophage colony-stimulating factor /GM-CSF, granulocyte colony-stimulating factor/G-CSF, IL-1 alpha, IL-1 beta, IL-6, IL-8 and TNF-alpha. Also interferes with antigen presentation by reducing the expression of MHC-class II and co-stimulatory molecules, thereby inhibiting their ability to induce T cell activation (By similarity). In addition, controls the inflammatory response of macrophages by reprogramming essential metabolic pathways including mTOR signaling (By similarity).

**Cellular Location**

Secreted {ECO:0000250 | UniProtKB:P22301}.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.