10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Shh

Catalog # PVGS1219

Product Information

Primary Accession	Q62226
Species	Mouse
Sequence	Cys25-Gly198(Cys25Ile-Val-Ile)
Purity	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
Endotoxin Level	
Expression System	E. coli
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH_2O up to 100 □g/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	20423
Other Names	Sonic hedgehog protein, SHH, 3.1.-.-, HHG-1, Shh unprocessed N-terminal signaling and C-terminal autoprocessing domains, ShhNC, Sonic hedgehog protein N-product, ShhN, Shh N-terminal processed signaling domains, ShhNp, Sonic hedgehog protein 19 kDa product, Shh {ECO:0000312|MGI:MGI:98297}, Hhg1
Target Background	Members of the Hedgehog (Hh) family are highly conserved proteins which are widely represented throughout the animal kingdom. The three known mammalian Hh proteins, Sonic (Shh), Desert (Dhh) and Indian (Ihh) are structurally related and share a high degree of amino-acid sequence identity (e.g., Shh and Ihh are 93% identical). The biologically active form of Hh molecules is obtained by autocatalytic cleavage of their precursor proteins and corresponds to approximately the N-terminal one half of the precursor molecule. Although Hh proteins have unique expression patterns and distinct biological roles within their respective regions of secretion, they use the same signaling pathway and can substitute for each other in experimental systems.

Protein Information

Name	Shh {ECO:0000303|PubMed:7916661, ECO:0000312|MGI:MGI:98297}
Function	[Sonic hedgehog protein]: Precursor of sonic hedgehog, a morphogen that activates the smoothened signaling pathway, and which is essential for a variety of patterning events during development (PubMed:11389830, PubMed:31657721, PubMed:7736596, PubMed:7891723, PubMed:8824192). The C-terminal part of the precursor displays an autoproteolysis and a cholesterol transferase activity, resulting (1) in the cleavage of the full-length protein into two parts, Sonic hedgehog protein N-product and C-product (ShhN and ShhC, respectively) and (2) covalent attachment of a cholesterol moiety to the C-terminus of the newly generated ShhN (PubMed:7736596, PubMed:7891723, PubMed:8824192). Both autoproteolysis and a cholesterol transferase activities occur in the endoplasmic reticulum (PubMed:21357747). Following additional lipidation, ShhN acts as a morphogen, while ShhC is degraded in the endoplasmic reticulum (PubMed:21357747).
Cellular Location	[Sonic hedgehog protein]: Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q15465}. Golgi apparatus membrane {ECO:0000250|UniProtKB:Q15465}. Note=Co-localizes with HHAT in the ER and Golgi membrane. {ECO:0000250|UniProtKB:Q15465}
Tissue Location	Expressed in a number of embryonic tissues including the notochord, ventral neural tube, floor plate, lung bud, zone of polarizing activity and posterior distal mesenchyme of limbs (PubMed:7916661). In the adult, expressed in lung and neural retina (PubMed:7916661).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.