

# IL-1 $\beta$

Catalog # PVGS1223

## Product Information

---

<b>Primary Accession Species</b>	<a href="#">P01584</a> Human
<b>Sequence</b>	Ala117-Ser269
<b>Purity</b>	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
<b>Endotoxin Level</b>	
<b>Expression System</b>	CHO
<b>Formulation</b>	Lyophilized after extensive dialysis against PBS.
<b>Reconstitution</b>	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH <sub>2</sub> O or PBS up to 100 $\mu$ g/ml.
<b>Storage &amp; Stability</b>	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

## Additional Information

---

<b>Gene ID</b>	3553
<b>Other Names</b>	Interleukin-1 beta, IL-1 beta, Catabolin, IL1B ( <a href="#">HGNC:5992</a> ), IL1F2
<b>Target Background</b>	Interleukin 1 beta is a proinflammatory cytokine produced in a variety of cells including monocytes, tissue macrophages, keratinocytes and other epithelial cells. Both IL-1 alpha and IL-1 beta binds to the same receptor and has similar if not identical biological properties. These cytokines have a broad range of activities including, stimulation of thymocyte proliferation, by inducing IL-2 release, B-cell maturation and proliferation, mitogenic FGF-like activity and the ability to stimulate the release of prostaglandin and collagenase from synovial cells. However, whereas IL-1 beta is a secreted cytokine, IL-1 alpha is predominantly a cell-associated cytokine.

## Protein Information

---

<b>Name</b>	IL1B ( <a href="#">HGNC:5992</a> )
-------------	------------------------------------

<b>Synonyms</b>	IL1F2
<b>Function</b>	Potent pro-inflammatory cytokine (PubMed: <a href="#">10653850</a> , PubMed: <a href="#">12794819</a> , PubMed: <a href="#">28331908</a> , PubMed: <a href="#">3920526</a> ). Initially discovered as the major endogenous pyrogen, induces prostaglandin synthesis, neutrophil influx and activation, T-cell activation and cytokine production, B-cell activation and antibody production, and fibroblast proliferation and collagen production (PubMed: <a href="#">3920526</a> ). Promotes Th17 differentiation of T-cells. Synergizes with IL12/interleukin-12 to induce IFNG synthesis from T-helper 1 (Th1) cells (PubMed: <a href="#">10653850</a> ). Plays a role in angiogenesis by inducing VEGF production synergistically with TNF and IL6 (PubMed: <a href="#">12794819</a> ). Involved in transduction of inflammation downstream of pyroptosis: its mature form is specifically released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed: <a href="#">33377178</a> , PubMed: <a href="#">33883744</a> ). Acts as a sensor of S.pyogenes infection in skin: cleaved and activated by pyogenes SpeB protease, leading to an inflammatory response that prevents bacterial growth during invasive skin infection (PubMed: <a href="#">28331908</a> ).
<b>Cellular Location</b>	Cytoplasm, cytosol. Secreted. Lysosome Secreted, extracellular exosome {ECO:0000250 UniProtKB:P10749} Note=The precursor is cytosolic (PubMed:15192144). In response to inflammasome-activating signals, such as ATP for NLRP3 inflammasome or bacterial flagellin for NLRC4 inflammasome, cleaved and secreted (PubMed:24201029, PubMed:33377178, PubMed:33883744). Mature form is secreted and released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed:33883744). In contrast, the precursor form is not released, due to the presence of an acidic region that is proteolytically removed by CASP1 during maturation (PubMed:33883744). The secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10 (PubMed:32272059)
<b>Tissue Location</b>	Expressed in activated monocytes/macrophages (at protein level).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.