

# GRO- $\alpha$ /MGSA/CXCL1

Catalog # PVGS1247

## Product Information

---

<b>Primary Accession</b>	<a href="#">P09341</a>
<b>Species</b>	Human
<b>Sequence</b>	Ala35-Asn107
<b>Purity</b>	> 95% as analyzed by SDS-PAGE
<b>Endotoxin Level</b>	
<b>Biological Activity</b>	Active, measured in a functional assay using HUVEC cells.
<b>Expression System</b>	CHO
<b>Formulation</b>	Lyophilized after extensive dialysis against PBS.
<b>Reconstitution</b>	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH <sub>2</sub> O or PBS up to 100 $\mu$ g/ml.
<b>Storage &amp; Stability</b>	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

## Additional Information

---

<b>Gene ID</b>	2919
<b>Other Names</b>	Growth-regulated alpha protein, C-X-C motif chemokine 1, GRO-alpha(1-73), Melanoma growth stimulatory activity, MGSA, Neutrophil-activating protein 3, NAP-3, GRO-alpha(4-73), GRO-alpha(5-73), GRO-alpha(6-73), CXCL1, GRO, GRO1, GROA, MGSA, SCYB1
<b>Target Background</b>	GRO- $\alpha$ /MGSA/CXCL1 has chemotactic activity for neutrophils. It may play a role in inflammation and exerts its effects on endothelial cells in an autocrine fashion. All three isoforms of GRO are CXC chemokines that can signal through the CXCR1 or CXCR2 receptors. GRO expression is inducible by serum or PDGF and/or by a variety of inflammatory mediators, such as IL-1 and TNF, in monocytes, fibroblasts, melanocytes and epithelial cells. In certain tumor cell lines, GRO is expressed constitutively.

## Protein Information

---

<b>Name</b>	CXCL1
<b>Synonyms</b>	GRO, GRO1, GROA, MGSA, SCYB1
<b>Function</b>	Has chemotactic activity for neutrophils. May play a role in inflammation and exerts its effects on endothelial cells in an autocrine fashion. In vitro, the processed forms GRO-alpha(4-73), GRO- alpha(5-73) and GRO-alpha(6-73) show a 30-fold higher chemotactic activity.
<b>Cellular Location</b>	Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.