10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



KGF/FGF-7

Catalog # PVGS1270

Product Information

Primary Accession P21781
Species Human

Sequence Cys32-Thr194

Purity > 95% as analyzed by SDS-PAGE

> 95% as analyzed by HPLC

Endotoxin Level

Expression System E. coli

Formulation Lyophilized after extensive dialysis against PBS.

Reconstitution It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

 ddH_2O up to 100 $\square g/ml$.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at lower than

-70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw

cycles.

Additional Information

Gene ID 2252

Other Names Fibroblast growth factor 7, FGF-7, Heparin-binding growth factor 7, HBGF-7,

Keratinocyte growth factor, FGF7, KGF

Target Background Keratinocyte Growth Factor (KGF) is a highly specific epithelial mitogen

produced by fibroblasts and mesenchymal stem cells. KGF belongs to the heparin binding Fibroblast Growth Factor (FGF) family, and is known as FGF-7. However, in contrast to the FGF-1, which binds to all known FGF receptors with high affinity, KGF only binds to a splice variant of an FGF receptor, FGFR2-IIIb. FGFR2-IIIb is produced by most of the epithelial cells, indicating that KGF plays roles as a paracrine mediator. KGF induces the differen-tiation and proliferation of various epithelial cells, including keratinocytes in the epidermis, hair follicles and sebaceous glands, and is responsible for the wound repairs of various tissues, including lung, bladder, and kidney.

Protein Information

Name FGF7

Synonyms KGF

Function Plays an important role in the regulation of embryonic development, cell

proliferation and cell differentiation. Required for normal branching morphogenesis. Growth factor active on keratinocytes. Possible major

paracrine effector of normal epithelial cell proliferation.

Cellular Location Secreted.

Tissue Location Epithelial cell.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.