

IGF-BP-4

Catalog # PVGS1314

Product Information

Primary Accession P22692
Species Human

Sequence Asp22-Glu258

Purity > 95% as analyzed by SDS-PAGE

Endotoxin Level

Biological Activity ED₅₀ **Expression System** HEK 293

Formulation Lyophilized after extensive dialysis against PBS.

Reconstitution It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

ddH₂O or PBS up to 100 □g/ml.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at -70°C or

-20°C. Upon reconstitution, the product should be stable for up to 1 week at

4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID 3487

Other Names Insulin-like growth factor-binding protein 4, IBP-4, IGF-binding protein 4,

IGFBP-4, IGFBP4, IBP4

Target Background Insulin-like growth factor-binding protein 4 (IGF-BP-4), also known as IBP-4, is

a secreted glycoprotein belonging to the IGFBP family. IGF-BP-4 is produced by osteoblasts, epidermis, ovarian follicles and other tissues. It binds both insulin-like growth factor (IGF) I and II, and it circulates in the plasma in both glycosylated and non-glycosylated forms. IGF-BP-4 prolongs the half-life of the IGFs and has been shown to inhibit or stimulate the growth-promoting effects of the IGFs. Pregnancy Associated Plasma Protein A (PAPP-A)

proteolytically cleaves IGF-BP-4 and reduces its affinity to bind IGFs, and thus

serves as an important regulator of IGF-BP-4 function.

Protein Information

Name IGFBP4

Synonyms IBP4

Function

IGF-binding proteins prolong the half-life of the IGFs and have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell

culture. They alter the interaction of IGFs with their cell surface receptors.

Cellular Location Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.