

NT-4

Catalog # PVGS1326

Product Information

Primary Accession Species	Q80VU4 Mouse
Sequence	Gly80-Ala209, expressed with an N-terminal Met
Purity	> 95% as analyzed by SDS-PAGE
Endotoxin Level	
Expression System	E. coli
Formulation	Lyophilized after extensive dialysis against 50 mM acetic acid.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in 50 mM acetic acid or ddH ₂ O up to 100 µg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	78405
Other Names	Neurotrophin-4, NT-4, Neurotrophin-5, NT-5, Neutrophic factor 4, Ntf4, Ntf5
Target Background	Neurotrophin-4 (NT-4), also known as NT-5, is a neurotrophic factor structurally related to β-NGF, BDNF, and NT-3. Human NT-4 shares 48 - 52% aa sequence identity with human β-NGF, BDNF, and NT-3. Neurotrophins have six conserved cysteine residues that are involved in the formation of three disulfide bonds. NT-4 is expressed highest levels in prostate, lower levels in thymus, placenta, and skeletal muscle. NT-4 binds and induces receptor dimerization and activation of TrkB. NT-4 can signal through TrkB receptors and promotes the survival of peripheral sensory sympathetic neurons.

Protein Information

Name	Ntf4
Synonyms	Ntf5

Function	Target-derived survival factor for peripheral sensory sympathetic neurons (By similarity). May promote ameloblast differentiation and subsequent reduction in proliferation of ameloblasts (PubMed: 27015268).
Cellular Location	Secreted {ECO:0000250 UniProtKB:P34130}.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.