

OSM

Catalog # PVGS1361

Product Information

Primary Accession	Q65Z15
Species	Rat
Sequence	MKRGCS SSP KLLSQLKSQA NITGNTASLL EPYILHQNLN TLTLRAACTE HPVAFPS EDM LRQLSKPDFL STVHATLGRV WHQLGA FRQQ FPKIQDFPEL ERARQNIQGI RNNVYCMARL LHPPLEIPEP TQADSGTSRP TTTAPGIFQI KIDSCRFLWG YHRFMGSVGR VFEEWGDGSR RSRRHSPLWA WLKGDHRIRP SRSSQSAMLRLSLVPR
Purity	> 95% as analyzed by SDS-PAGE and HPLC.
Endotoxin Level	
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O or PBS at 100 µg/ml.

Additional Information

Gene ID	289747
Other Names	Oncostatin-M, OSM, Osm
Target Background	<p>Oncostatin M (OSM) is a multifunctional cytokine, and belongs to Interleukin-6 (IL-6) subfamily, which also includes IL-11, leukemia inhibitory factor (LIF), ciliary neurotropic factor, cardiotrophin-1, and novel neurotrophin-1. In vivo, OSM is secreted from activated T cells, monocytes, neutrophils, and endothelial cells. OSM is related to LIF, and shares a receptor with LIF in human. Human OSM can bind to gp130 and recruit OSM Receptor β or LIF Receptor β to form a ternary complex. OSM stimulates the growth of different types of cells, including megakaryocytes, fibroblasts, vascular endothelial cells, and T cells. OSM inhibits the proliferation of several cancer cell lines, such as solid tissue tumor cells, lung cancer cells, melanoma cells, and breast cancer cells.</p> <p>Recombinant Rat Oncostatin M (rrOSM) produced in E. coli is a single non-glycosylated polypeptide chain containing 215 amino acids. A fully biologically active molecule, rrOSM has a molecular mass of 24.5 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at .</p>

Protein Information

Name	Osm
-------------	-----

Function	Growth regulator. Inhibits the proliferation of a number of tumor cell lines. It regulates cytokine production, including IL-6, G-CSF and GM-CSF from endothelial cells (By similarity). Uses only type II OSM receptor (heterodimers composed of OSMR and IL6ST). Involved in the maturation of fetal hepatocytes, thereby promoting liver development and regeneration.
Cellular Location	Secreted.
Tissue Location	Widely expressed. Expressed at higher levels in liver, skin and spleen.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.