

ORFX, RING3L

Catalog # PVGS1374

Product Information

Primary Accession NM_007371
Species Human

Sequence MHHHHHHPSK PGRKTNQLQY MQNVVVKTLW KHQFAWPFYQ PVDAIKLNLP

DYHKIIKNPM DMGTIKKRLE NNYYWSASEC MQDFNTMFTN CYIYNKPTDD

IVLMAQALEK IFLQKVAQMP QEEV

Purity > 95% by SDS-PAGE and HPLC analysis.

Endotoxin Level Formulation

Sterile liquid solution contains 25mM HEPES, pH7.5, 150mM NaCl, 5%

glycerol, 0.5 mM TCEP. Frozen solution.

Additional Information

Target Background

Bromodomain (BRD) is an extensive family of protein domains, originally identified in and named after the Drosophila protein Brahma. Members of BRD family share a conserved atypical left-handed four helix bundle structure, and specifically bind to ε-lysine acetylated proteins. It is well known that histone acetylation and methylation play a central role in epigenetics and are important for various gene transcription events, thus the acetyl-lysine binding property of BRDs make them suitable drug targets for epigenetics. Currently, there are 46 diverse human proteins containing 61 BRDs. These include histone acetyltransferases, ATP-dependent chromatin-remodeling complex proteins, and nuclear scaffold proteins. The main functions of BRDs in vivo include chromatin acetylation and deacetylation, nucleosome assembly and remodeling, and organizations of chromosome or chromatin domains. Recombinant human BRD3 (29-145) with His tag produced in E. coli is a single, non-glycosylated polypeptide chain containing 124 amino acids. A fully biologically active molecule, BRD3 (29-145) has a molecular mass of 14.9 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at .

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.