

# SCF

Catalog # PVGS1379

## Product Information

---

|                                  |  |
|----------------------------------|--|
| <b>Primary Accession Species</b> | <a href="#">P21581</a><br>Rat  |
| <b>Sequence</b>                  | Gln26-Ala189   |
| <b>Purity</b>                    | > 95% as analyzed by SDS-PAGE  |
| <b>Endotoxin Level</b>           |  |
| <b>Biological Activity</b>       | ED <sub>50</sub>   |
| <b>Expression System</b>         | HEK 293  |
| <b>Formulation</b>               | Lyophilized after extensive dialysis against PBS.  |
| <b>Reconstitution</b>            | It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH <sub>2</sub> O or PBS up to 100 µg/ml.  |
| <b>Storage &amp; Stability</b>   | Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles. |

## Additional Information

---

|                          |   |
|--------------------------|---|
| <b>Gene ID</b>           | 60427   |
| <b>Other Names</b>       | Kit ligand, Hematopoietic growth factor KL, Mast cell growth factor, MGF, Stem cell factor, SCF, c-Kit ligand, Soluble KIT ligand, sKITLG, Kitlg, Kitl, Mgf   |
| <b>Target Background</b> | Stem cell factor (also known as SCF, KIT-ligand, KL, or steel factor) is a cytokine that binds to the c-KIT receptor (CD117). SCF can exist both as a transmembrane protein and a soluble protein. It stimulates the proliferation of myeloid, erythroid, and lymphoid progenitors in bone marrow cultures and has been shown to act synergistically with colony stimulating factors. SCF plays an important role in the hematopoiesis during embryonic development. SCF can regulate HSCs in the stem cell niche in the bone marrow. SCF has been shown to increase the survival of HSCs in vitro and contributes to the self-renewal and maintenance of HSCs in-vivo. |

## Protein Information

---

|             |       |
|-------------|-------|
| <b>Name</b> | Kitlg |
|-------------|-------|



|                          |   |
|--------------------------|---|
| <b>Synonyms</b>          | Kitl, Mgf   |
| <b>Function</b>          | <p>Ligand for the receptor-type protein-tyrosine kinase KIT. Plays an essential role in the regulation of cell survival and proliferation, hematopoiesis, stem cell maintenance, gametogenesis, mast cell development, migration and function, and in melanogenesis. KITLG/SCF binding can activate several signaling pathways. Promotes phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, and subsequent activation of the kinase AKT1. KITLG/SCF and KIT also transmit signals via GRB2 and activation of RAS, RAF1 and the MAP kinases MAPK1/ERK2 and/or MAPK3/ERK1. KITLG/SCF and KIT promote activation of STAT family members STAT1, STAT3 and STAT5. KITLG/SCF and KIT promote activation of PLCG1, leading to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. KITLG/SCF acts synergistically with other cytokines, probably interleukins.</p> |
| <b>Cellular Location</b> | [Isoform 1]: Cell membrane; Single- pass type I membrane protein [Soluble KIT ligand]: Secreted.  |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.