

HCGF

Catalog # PVGS1380

Product Information

Primary Accession	P97688
Species	Rat
Sequence	MISDRGSDAH HLLRTLDCRT IALEILVKLP YPQVSGLNNS DDKANLRNST LRRVNLDEFL KSQEEFDSQD TTDIKSKLQK LKCCIPAAAS DSVLPGVYNK DLDDFKKKLR FYVIHLKDLQ PVSVSRRPPQP TSSSDNFRPM TVEC
Purity	> 95% as analyzed by SDS-PAGE and HPLC.
Endotoxin Level	
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O at 100 µg/mL.

Additional Information

Target Background	<p>Interleukin-3 (IL-3) is a pleiotropic cytokine belonging to the interleukin family. IL-3 shares similarities with Granulocyte-Macrophage Colony-Stimulating Factor (GM-CSF) and IL-5: they all have a four-helix bundle structure, are located on the same chromosomes in both human and mouse, are produced by activated T cells, and share receptors. The IL-3/IL-5/GM-CSF receptor family members are all heterodimeric, composed of a receptor-specific α chain and a common β chain. IL-3 is also called multi-colony stimulating factor since it stimulates the development and colony formation of multiple lineages of hematopoietic cells by activating intracellular pathways such as Ras-Raf-ERK and JAK/STAT. IL-3 inhibits apoptosis and promotes cell survival by targeting the anti-apoptotic bcl-2 gene family.</p> <p>Recombinant rat Interleukin-3 beta (rrIL-3 beta) produced in E. coli is a single non-glycosylated polypeptide chain containing of 144 amino acids. A fully biologically active molecule, rrIL-3 beta has a molecular mass of 16.3 kDa analyzed by non-reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at .</p>
--------------------------	---

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.