

CXCL3, Human

Catalog # PVGS1384

Product Information

Primary Accession	P19876
Species	Human
Sequence	ASVVTELRCQ CLQTLQGIHL KNIQSVNVRS PGPHCAQTEV IATLKNGKKA CLNPASPMVQ KIEKILNKG STN
Purity	> 98% as analyzed by SDS-PAGE.
Endotoxin Level	
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O or PBS at 100 µg/ml.

Additional Information

Gene ID	2921
Other Names	C-X-C motif chemokine 3, GRO-gamma(1-73), Growth-regulated protein gamma, GRO-gamma, Macrophage inflammatory protein 2-beta, MIP2-beta, GRO-gamma(5-73), CXCL3, GRO3, GROG, SCYB3
Target Background	<p>Chemokine (C-X-C motif) ligand 3 (CXCL3) is a small cytokine belonging to the CXC chemokine family that is also known as GRO3 oncogene (GRO3), GRO protein gamma (GROg) and macrophage inflammatory protein-2-beta (MIP2b). CXCL3 controls migration and adhesion of monocytes and mediates its effect on its target cell by interacting with cell surface chemokine receptor CXCR2. It has been shown that CXCL3 regulates the migration of precursors of cerebellar granule neurons toward the internal layers of the cerebellum, during morphogenesis.</p> <p>Recombinant humanGRO-gamma/CXCL3 produced in CHO cells is a polypeptide chain containing 73 amino acids. A fully biologically active molecule, rhGRO-gamma/CXCL3 has a molecular mass of 9kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at .</p>

Protein Information

Name	CXCL3
Synonyms	GRO3, GROG, SCYB3
Function	Ligand for CXCR2 (By similarity). Has chemotactic activity for neutrophils. May play a role in inflammation and exert its effects on endothelial cells in an autocrine fashion. In vitro, the processed form GRO-gamma(5-73) shows a

fivefold higher chemotactic activity for neutrophilic granulocytes.

Cellular Location

Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.