10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

LIX/CXCL5

Catalog # PVGS1386

Product Information

Primary Accession	P50228
Species	Mouse
Sequence	Val45-Ala118
Purity	> 98% as analyzed by SDS-PAGE
Endotoxin Level	
Biological Activity	The EC_{50} value of mouse LIX/CXCL5 (74aa) on Ca^{2+} mobilization assay in CHO-K1/Gα15/mCXCR2 cells (human Gα15 and mouse CXCR2 stably expressed in CHO-K1 cells) is less than 200.0 ng/ml.
Expression System	HEK 293 Cells
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH_2O or PBS up to 100 □g/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	20311
Other Names	C-X-C motif chemokine 5, Cytokine LIX, Small-inducible cytokine B5, GCP-2(1-78), GCP-2(9-78), Cxcl5, Scyb5
Target Background	Mouse LIX (C-X-C motif chemokine 5) is a small cytokine belonging to the CXC chemokine family that is cleaved into the following 2 chains [GCP-2(1-78) and GCP-2(9-78)]. Mouse LIX plays a role in reducing sensitivity to sunburn pain in some subjects, and is a potential target which could be used to understand more about pain in other inflammatory conditions. It is most closely related to two highly homologous human neutrophil chemoattractants GCP-2 and ENA-78. The first 78 amino acid residues within the predicted mature mouse LIX shares approximately 61% and 55% amino acid identity with human GCP-2 and ENA-78. This chemokine stimulates the chemotaxis of neutrophils possessing angiogenic properties. It elicits these effects by interacting with the cell surface chemokine receptor CXCR2.

Protein Information

Name	Cxcl5
Synonyms	Scyb5
Function	May participate in the recruitment of inflammatory cells by injured or infected tissue. GCP-2(1-78) and, more potent, GCP-2(9-78) attract neutrophils and are involved in neutrophil activation.
Cellular Location	Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.