10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

LD78-β/CCL3L1

Catalog # PVGS1387

Product Information

Primary Accession	P16619
Species	Human
Sequence	Ala24-Ala93
Purity	> 95% as analyzed by SDS-PAGE > 95% as analyzed by HPLC
Endotoxin Level	
Expression System	E. coli
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH_2O up to 100 □g/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	414062;6349
Other Names	C-C motif chemokine 3-like 1, G0/G1 switch regulatory protein 19-2, LD78-beta(1-70), PAT 464.2, Small-inducible cytokine A3-like 1, Tonsillar lymphocyte LD78 beta protein, LD78-beta(3-70), LD78-beta(5-70), CCL3L1, D17S1718, G0S19-2, SCYA3L1
Target Background	LD78-beta/CCL3L1 is a proinflammatory chemokine and the isoform of Macrophage Inflammatory Protein-1 alpha (MIP-1 alpha). LD78-beta is secreted by most mature leukocytes, predominantly macrophages, and its major receptor is the G-protein coupled receptor CCR5, which is also the co-receptor used by the HIV-1 virus for cell entry. LD78-beta has superior antiviral activity and induces a variety of immune cells, particularly $CD8^+$> T cells and immature dendritic cells. LD78-beta attracts lymphocytes and macrophages to sites of inflammation and infection, and its functions are inhibited by Interleukin-4, Interleukin-10, and Interleukin-13. Importantly, the copy number variation of LD78-beta is associated with HIV susceptibility, indicating LD78-beta's critical role in the disease.

Protein Information

Name	CCL3L1
Synonyms	D17S1718, G0S19-2, SCYA3L1
Function	Chemotactic for lymphocytes and monocytes. Is a ligand for CCR1, CCR3 and CCR5. Is an inhibitor of HIV-1-infection. The processed form LD78-beta(3-70) shows a 20-fold to 30-fold higher chemotactic activity and is a very potent inhibitor of HIV-1-infection. LD78- beta(3-70) is also a ligand for CCR1, CCR3 and CCR5.
Cellular Location	Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.