10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ENA-78/CXCL5

Catalog # PVGS1389

Product Information

Primary Accession	P42830
Species	Human
Sequence	Ala37-Asn114
Purity	> 98% as analyzed by SDS-PAGE
Endotoxin Level	
Biological Activity	The EC_{50} value of human ENA-78/CXCL5 on Ca^{2+} mobilization assay in CHO-K1/Gα15/hCXCR2 cells (human Gα15 and human CXCR2 stably expressed in CHO-K1 cells) is less than 200.0 ng/ml.
Expression System	HEK 293
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH_2O or PBS up to 100 □g/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	6374
Other Names	C-X-C motif chemokine 5, ENA-78(1-78), Epithelial-derived neutrophil-activating protein 78, Neutrophil-activating peptide ENA-78, Small-inducible cytokine B5, ENA-78(8-78), ENA-78(9-78), CXCL5, ENA78, SCYB5
Target Background	Epithelial cellderived neutrophilactivating peptide (ENA78) also known as C-X-C motif chemokine 5 (CXCL5), is a small cytokine belonging to the CXC chemokine family. It is produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor-alpha. Expression of CXCL5 has also been observed in eosinophils, and can be inhibited with the type II interferon, IFN-γ. This chemokine stimulates the chemotaxis of neutrophils possessing angiogenic properties. Full length CXCL5 (78 aa) is trimmed at the Nterminal end by cathepsin G and chymotrypsin to ENA-74 (74 aa) and ENA-70 (70aa), with the shortened forms showing increased potency relative to full length CXCL5. CXCL5can signal

through the CXCR2 receptor.

Protein Information

Name	CXCL5
Synonyms	ENA78, SCYB5
Function	Involved in neutrophil activation. In vitro, ENA-78(8-78) and ENA-78(9-78) show a threefold higher chemotactic activity for neutrophil granulocytes.
Cellular Location	Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.