

# CXCL5, ENA-78

Catalog # PVGS1394

## Product Information

---

<b>Primary Accession</b>	<a href="#">P97885.1</a>
<b>Species</b>	Rat
<b>Sequence</b>	APFSAMVATE LRCVCLTLAP RINPKMIANL EVIPAGPHCP KVEVIAKLKN QKDNVCLDPQ APLIKKVIQK ILGSENKKTK RNALALVR
<b>Purity</b>	> 98% as analyzed by SDS-PAGE.
<b>Endotoxin Level</b>	
<b>Formulation</b>	Lyophilized after extensive dialysis against PBS.
<b>Reconstitution</b>	Reconstituted in ddH <sub>2</sub> O or PBS at 100 µg/ml.

## Additional Information

---

<b>Target Background</b>	<p>LPSinduced CXC chemokine (LIX), also known as C-X-C motif chemokine 5(CXCL5), is a small cytokine belonging to the CXC chemokine family that is also known as epithelial-derived neutrophil-activating peptide 78 (ENA-78). It is produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor-alpha. Rat LIX cDNA encodes a 130 aa residue precursor with a predicted 37 aa residue signal peptide and a 93 aa residue mature protein. Among human CXC chemokines, rat LIX is most closely related to human GCP-2 and ENA-78. LIX can signal through the CXCR2 receptor.</p> <p>Recombinant rat LIX/CXCL5 (88aa) produced in CHO cells is a polypeptide chain containing 88 amino acids. A fully biologically active molecule, rrLIX/CXCL5 (88aa) has a molecular mass of 9.6 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at .</p>
--------------------------	--

## Protein Information

---

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.