

MIP-2/CXCL2 Catalog # PVGS1401

Product Information

Primary Accession P10889
Species Mouse

Sequence Gly27-Asn100

Purity > 95% as analyzed by SDS-PAGE

Endotoxin Level

Biological Activity The EC₅₀ value of mouse MIP-2/CXCL2 on Ca²⁺ mobilization assay in

CHO-K1/Ga15/mCXCR2 cells (human Ga15 and mouse CXCR2 stably

expressed in CHO-K1 cells) is less than 1.0 ng/ml.

Expression System E. coli

Formulation Lyophilized after extensive dialysis against PBS.

Reconstitution It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

ddH₂O or PBS up to 100 □g/ml.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at lower than

-70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw

cycles.

Additional Information

Gene ID 20310

Other Names C-X-C motif chemokine 2, Macrophage inflammatory protein 2, MIP2, Cxcl2,

Mip-2, Mip2, Scyb2

Target Background Chemokine (C-X-C motif) ligand 2 (CXCL2) is a small cytokine belonging to the

CXC chemokine family that is also referred to as macrophage inflammatory protein 2-alpha (MIP2-alpha), Growth-regulated protein beta (Gro-beta) and Gro oncogene-2 (Gro-2). CXCL2 is secreted by monocytes and macrophages and is chemotactic for polymorphonuclear leukocytes and hematopoietic stem cells. CXCL2's amino acid sequence is 90% identical to the amino acid sequence of related chemokine, CXCL1. CXCL2 signals through the CXCR2

receptor.

Protein Information

Name Cxcl2

Synonyms Mip-2, Mip2, Scyb2

Chemotactic for human polymorphonuclear leukocytes but does not induce chemokinesis or an oxidative burst. **Function**

Secreted. **Cellular Location**

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.