10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HCC-1/CCL14

Catalog # PVGS1406

Product Information

Primary Accession	Q16627
Species	Human
Sequence	Thr22-Asn93
Purity	> 95% as analyzed by SDS-PAGE
Endotoxin Level	
Expression System	E. coli
Theoretical Molecular Weight	8.4 kDa
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH_2O up to 100 □g/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	6358
Other Names	C-C motif chemokine 14, Chemokine CC-1/CC-3, HCC-1/HCC-3, HCC-1(1-74), NCC-2, Small-inducible cytokine A14, HCC-1(3-74), HCC-1(4-74), HCC-1(9-74), CCL14, NCC2, SCYA14
Target Background	HCC-1/CCL14 is a member of the chemokine family, which are small chemotactic proteins that regulate cell migration under inflammatory and steady state conditions. HCC-1 is expressed in epithelial and decidual cells and is unique among chemokines due to its high abundance in normal human plasma. HCC-1 can bind to chemokine receptors CCR1 and CCR5, however full length HCC-1 is a weak agonist of CCR1 and only becomes potent after removal of its eight N-terminal residues. Chemokine decoy receptor D6 can bind HCC-1 and promote its degradation as a means to regulate its level in vivo. Functionally HCC-1 promotes trophoblast migration by regulating extracellular matrix components as well as specific adhesion molecules.

Protein Information

Name	CCL14
Synonyms	NCC2, SCYA14
Function	Has weak activities on human monocytes and acts via receptors that also recognize MIP-1 alpha. It induces intracellular Ca(2+) changes and enzyme release, but no chemotaxis, at concentrations of 100-1,000 nM, and is inactive on T-lymphocytes, neutrophils, and eosinophil leukocytes. Enhances the proliferation of CD34 myeloid progenitor cells. The processed form HCC-1(9-74) is a chemotactic factor that attracts monocytes, eosinophils, and T-cells and is a ligand for CCR1, CCR3 and CCR5.
Cellular Location	Secreted.
Tissue Location	Expressed constitutively in several normal tissues: spleen, liver, skeletal and heart muscle, gut, and bone marrow, present at high concentrations (1-80 nM) in plasma

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.