

## Noggin

Catalog # PVGS1416

## **Product Information**

Primary Accession P97466
Species Mouse

Sequence Leu20-Cys232

**Purity** > 95% as analyzed by SDS-PAGE

**Endotoxin Level** 

**Biological Activity** ED<sub>50</sub> Expression System CHO

**Formulation** Lyophilized after extensive dialysis against PBS.

**Reconstitution** It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

ddH<sub>2</sub>O or PBS up to 100 □g/ml.

**Storage & Stability** Upon receiving, this product remains stable for up to 6 months at lower than

-70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw

cycles.

## **Additional Information**

**Gene ID** 18121

Other Names Noggin, Nog

**Target Background** Noggin, also known as NOG, is a homodimeric glycoprotein that bindsto and

modulates the activity of TGF-beta family ligands. It is expressed in

condensing cartilage and immature chondrocytes. Noggin antagonizes bone morphogenetic protein (BMP) activities by blocking epitopes on BMPs needed for binding to their receptors. Noggin has been shown to be involved in many developmental processes, such as neural tube formation and joint formation. During development, Noggin diffuses through extracellular matrices and forms morphogenic gradients, regulating cellular responses dependent on the

local concentration of the signaling molecule.

## **Protein Information**

Name Nog

**Function** Essential for cartilage morphogenesis and joint formation. Inhibitor of bone

morphogenetic proteins (BMP) signaling which is required for growth and

patterning of the neural tube and somite (PubMed: 9585504,

PubMed: 9603738). Inhibits chondrocyte differentiation through its interaction

with GDF5 and, probably, GDF6 (By similarity).

**Cellular Location** Secreted.

**Tissue Location** Expressed in condensing cartilage and immature chondrocytes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.