

FDP, MIAL1

Catalog # PVGS1424

Product Information

Sequence	Val ¹⁸ -Glu ¹²⁸ (Accession #: Q9NRC9)
Purity	> 95% as analyzed by SDS-PAGE.
Endotoxin Level	
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O or PBS at 100 µg/ml.

Additional Information

Target Background	<p>OTOR, also called Otoraplin and MIAL, is a secreted cytokine and a member of the melanoma-inhibiting activity gene family. Members of this family which also includes MIA, MIA2, and TANGO share a SRC homology-3 (SH3)-like domain. OTOR appears to be involved in early chondrogenesis of the otic capsule, which is required for normal inner ear development and auditory function. OTOR is highly homologous to MIA/cartilage-derived retinoic acid-sensitive protein (CD-RAP), which is a cartilage-specific protein that is also expressed in malignant melanoma cells. The 111 amino acid mature human otoraplin contains 1 SH3 domain (46-107 amino acids) and a Tyr at position 50 that is reportedly sulfated. Otoraplin takes part in the initiation of periotic mesenchyme chondrogenesis.</p> <p>Recombinant human Otoraplin (OTOR) produced in CHO cells is a single non-glycosylated polypeptide chain containing 111 amino acids. A fully biologically active molecule, rhOTOR has a molecular mass of 14-15 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at .</p>
--------------------------	---

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.