

I-TAC/CXCL11

Catalog # PVGS1437

Product Information

Primary Accession Species	O14625 Human
Sequence	Phe22-Phe94
Purity	> 98% as analyzed by SDS-PAGE
Endotoxin Level Biological Activity	The EC ₅₀ value of human I-TAC/CXCL11 on Ca ²⁺ mobilization assay in CHO-K1/Ga15/hCXCR3 cells (human Ga15 and human CXCR3 stably expressed in CHO-K1 cells) is less than 0.5 µg/ml.
Expression System	HEK 293
Formulation Reconstitution	Lyophilized after extensive dialysis against PBS. It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 µg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	6373
Other Names	C-X-C motif chemokine 11, Beta-R1, H174, Interferon gamma-inducible protein 9, IP-9, Interferon-inducible T-cell alpha chemoattractant, I-TAC, Small-inducible cytokine B11, CXCL11, ITAC, SCYB11, SCYB9B
Target Background	Chemokine (C-X-C motif) ligand 11(CXCL11), also known as I-TAC and B-R1, is a small cytokine belonging to the CXC chemokine family that is also called Interferon-inducible T-cell alpha chemoattractant (I-TAC) and Interferon-gamma-inducible protein 9 (IP-9). This chemokine elicits its effects on target cells by interacting with chemokine receptor CXCR3 having a higher affinity than other ligands for this receptor such as CXCL9 and CXCL10. CXCL11 is chemotactic for activated T cells. The gene encoding CXCL11 has been mapped to chromosome 4. CXCL11 cDNA encodes a 94 amino acid residue precursor protein with a 21 amino acid residue putative signal sequence, which is cleaved to form the mature 73 amino acid residue protein. CXCL11 shares 36% and 37% amino acid sequence homology with IP-10 and

MIG (two other known human non-ELR CXC chemokines), respectively. Mouse CXCL11 exhibits 68% sequence homology with human CXCL11.

Protein Information

Name	CXCL11
Synonyms	ITAC, SCYB11, SCYB9B
Function	Chemotactic for interleukin-activated T-cells but not unstimulated T-cells, neutrophils or monocytes. Induces calcium release in activated T-cells. Binds to CXCR3. May play an important role in CNS diseases which involve T-cell recruitment. May play a role in skin immune responses.
Cellular Location	Secreted.
Tissue Location	High levels in peripheral blood leukocytes, pancreas and liver astrocytes. Moderate levels in thymus, spleen and lung. Low levels in placenta, prostate and small intestine. Also found in epidermal basal layer keratinocytes in skin disorders

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.