10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

I-309/CCL1

Catalog # PVGS1455

Product Information

Primary Accession	P22362-1
Species	Human
Sequence	Lys24-Lys96
Purity	> 98% as analyzed by SDS-PAGE
Endotoxin Level	
Biological Activity	The EC_{50} value of human I-309/CCL1 on Ca^{2+} mobilization assay in CHO-K1/G15/hCCR8 cells (human G15 and human CCR8 stably expressed in CHO-K1 cells) is less than 1.0 □g/ml.
Expression System	CHO
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH_2O or PBS up to 100 □g/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Target Background	Chemokine (C-C motif) ligand 1 (CCL1), also known as I-309, is a small glycoprotein secreted by activated T cells that belongs to the family of chemokines. Human CCL1 has been assumed to be a homologue of mouse TCA3. While the two proteins share only approximately 42% amino acid sequence identity, both chemokines contain an extra pair of cysteine residues not found in most other chemokines. CCL1 attracts monocytes, NK cells, immature B cells and dendritic cells by interacting with the cell surface chemokine receptor CCR8. This chemokine resides in a large cluster of CC chemokines on human chromosome 17.

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.