

# I-309/CCL1

Catalog # PVGS1463

## Product Information

---

<b>Primary Accession Species</b>	<a href="#">P22362-1</a> Human
<b>Sequence</b>	Ser23-Lys96
<b>Purity</b>	> 95% as analyzed by SDS-PAGE
<b>Endotoxin Level Biological Activity</b>	The EC <sub>50</sub> value of human I-309/CCL on Ca <sup>2+</sup> mobilization assay in CHO-K1/Gα15/hCCR8 cells (human Gα15 and human CCR8 stably expressed in CHO-K1 cells) is less than 1.0 µg/ml.
<b>Expression System</b>	E. coli
<b>Theoretical Molecular Weight</b>	8.6 kDa
<b>Formulation Reconstitution</b>	Lyophilized after extensive dialysis against PBS. It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH <sub>2</sub> O or PBS up to 100 µg/ml.
<b>Storage &amp; Stability</b>	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

## Additional Information

---

<b>Target Background</b>	Chemokine (C-C motif) ligand 1 (CCL1), also known as I-309, is a small glycoprotein secreted by activated T cells that belongs to the family of chemokines. Human CCL1 has been assumed to be a homologue of mouse TCA3. While the two proteins share only approximately 42% amino acid sequence identity, both chemokines contain an extra pair of cysteine residues not found in most other chemokines. CCL1 attracts monocytes, NK cells, immature B cells and dendritic cells by interacting with the cell surface chemokine receptor CCR8. This chemokine resides in a large cluster of CC chemokines on human chromosome 17.
--------------------------	--

## Protein Information

---

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.