

MDC/CCL22

Catalog # PVGS1473

Product Information

Primary Accession 000626-1
Species Human

Sequence Pro26-Gln93

Purity > 95% as analyzed by SDS-PAGE

Endotoxin Level

Biological Activity The EC₅₀ value of human MDC/CCL22 (67aa) on Ca²⁺ mobilization assay in

CHO-K1/Gα15/hCCR4cells (human Gα15 and human CCR4 stably expressed in

CHO-K1 cells) is less than 1.0 [g/ml.

Expression System E. coli

Formulation Lyophilized after extensive dialysis against PBS.

Reconstitution It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

ddH₂O or PBS up to 100 □g/ml.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at lower than

-70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw

cycles.

Additional Information

Target Background

Macrophage-Derived/CCL22 Chemokine (MDC), also known as stimulated T cell chemotactic protein (STCP1), is a CC chemokine initially isolated from clones of monocytederived macrophages. CCL22 is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. CCL22 shows

chemotactic activity for natural killer cells, chronically activated T

lymphocytes, monocytes and dendritic cells. CCL22 has mild chemotactic activity for primary activated T lymphocytes and no chemoattractant activity for neutrophils, eosinophils or resting T lymphocytes. CCL22 may also be involved in certain aspects of activated T lymphocyte physiology, such astrafficking activated T lymphocytes to inflammatory sites. CCL22 interacts

with the cell surface chemokine receptor CCR4.

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.