

FGF-10

Catalog # PVGS1475

Product Information

Primary Accession O15520
Species Human

Sequence Leu40-Ser208

Purity > 95% as analyzed by SDS-PAGE

Endotoxin Level

Expression System E. coli

Formulation Lyophilized after extensive dialysis against PBS.

ReconstitutionIt is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

 ddH_2O up to 100 $\square g/ml$.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at lower than

-70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw

cycles.

Additional Information

Gene ID 2255

Other Names Fibroblast growth factor 10, FGF-10, Keratinocyte growth factor 2, FGF10

Target Background Fibroblast Growth Factor-10 (FGF-10) is a mitogen mainly produced by

mesenchymal stem cells in the lung. FGF-10 belongs to the heparin binding FGF family, and is also known as Keratinocyte Growth Factor-2 (KGF-2). It shares homology with KGF and receptor binding to FGFR2-IIIb. However, while KGF induces proliferation and differentiation of various epithelial cells,

FGF-10 promotes budding and branching morphogenesis during the

multi-organ development via mesenchymal-epithelial cell interactions. FGF-10 is critical for lung and limb development, and is regulated by Shh during early

development.

Protein Information

Name FGF10

Function Plays an important role in the regulation of embryonic development, cell

proliferation and cell differentiation. Required for normal branching morphogenesis. May play a role in wound healing.

Cellular Location

Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.