

MIP-2/CXCL2

Catalog # PVGS1484

Product Information

Primary Accession Species	P10889 Mouse
Sequence	Ala28-Asn100
Purity	> 95% as analyzed by SDS-PAGE
Endotoxin Level Biological Activity	The EC ₅₀ value of Mouse MIP-2/CXCL2 on Ca ²⁺ mobilization assay in CHO-K1/mCXCR2 (mouse CXCR2 stably expressed in CHO-K1 cells) is less than 200.0 ng/ml.
Expression System	HEK 293
Formulation Reconstitution	Lyophilized from a 0.2 µm filtered solution in PBS. It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 µg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	20310
Other Names	C-X-C motif chemokine 2, Macrophage inflammatory protein 2, MIP2, Cxcl2, Mip-2, Mip2, Scyb2
Target Background	Chemokine (C-X-C motif) ligand 2 (CXCL2) is a small cytokine belonging to the CXC chemokine family that is also referred to as macrophage inflammatory protein 2-alpha (MIP2-alpha), Growth-regulated protein beta (Gro-beta) and Gro oncogene-2 (Gro-2). CXCL2 is secreted by monocytes and macrophages and is chemotactic for polymorphonuclear leukocytes and hematopoietic stem cells. CXCL2's amino acid sequence is 90% identical to the amino acid sequence of related chemokine, CXCL1. CXCL2 signals through the CXCR2 receptor.

Protein Information

Name	Cxcl2
Synonyms	Mip-2, Mip2, Scyb2
Function	Chemotactic for human polymorphonuclear leukocytes but does not induce chemokinesis or an oxidative burst.
Cellular Location	Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.