

ENA-78/CXCL5

Catalog # PVGS1505

Product Information

Primary Accession Species	P42830 Human
Sequence	Arg45-Asn114
Purity	> 95% as analyzed by SDS-PAGE
Endotoxin Level Biological Activity	The EC ₅₀ value of human ENA78/CXCL5 (9-78 a.a.) on Ca ²⁺ mobilization assay in CHO-K1/G15/hCXCR2 cells (human G15 and human CXCR2 stably expressed in CHO-K1 cells) is less than 50.0 ng/ml.
Expression System	E. coli
Formulation Reconstitution	Lyophilized after extensive dialysis against PBS. It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 µg/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	6374
Other Names	C-X-C motif chemokine 5, ENA-78(1-78), Epithelial-derived neutrophil-activating protein 78, Neutrophil-activating peptide ENA-78, Small-inducible cytokine B5, ENA-78(8-78), ENA-78(9-78), CXCL5, ENA78, SCYB5
Target Background	Epithelial cell derived neutrophil activating peptide (ENA78) also known as C-X-C motif chemokine 5(CXCL5), is a small cytokine belonging to the CXC chemokine family. It is produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor-alpha. Expression of CXCL5 has also been observed in eosinophils, and can be inhibited with the type II interferon, IFN-γ. This chemokine stimulates the chemotaxis of neutrophils possessing angiogenic properties. Full length CXCL5 (78 aa) is trimmed at the Nterminal end by cathepsin G and chymotrypsin to ENA-74 (74 aa) and ENA-70 (70aa), with the shortened forms showing increased potency relative to full length CXCL5. CXCL5 can signal

through the CXCR2 receptor.

Protein Information

Name	CXCL5
Synonyms	ENA78, SCYB5
Function	Involved in neutrophil activation. In vitro, ENA-78(8-78) and ENA-78(9-78) show a threefold higher chemotactic activity for neutrophil granulocytes.
Cellular Location	Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.