

IP-10/CXCL10

Catalog # PVGS1507

Product Information

Primary Accession Q548V9
Species Mouse

Sequence Ile22-Pro98

Purity > 95% as analyzed by SDS-PAGE

Endotoxin Level

Biological Activity The EC₅₀ value of Mouse IP-10 on Ca²⁺ mobilization assay in

CHO-K1/Ga15/mCXCR3 cells (human Ga15 and mouse CXCR3 stably

expressed in CHO-K1 cells) is less than 2.0 [g/ml.

Expression System E. coli

Formulation Lyophilized after extensive dialysis against PBS.

Reconstitution It is recommended that this vial be briefly centrifuged prior to opening to

bring the contents to the bottom. Reconstitute the lyophilized powder in

ddH₂O or PBS up to 100 □g/ml.

Storage & Stability Upon receiving, this product remains stable for up to 6 months at lower than

-70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw

cycles.

Additional Information

Target Background

Interferon Inducible Protein 10 (IP-10)/CXCL10 was originally identified as an IFN-γ-inducible gene in monocytes, fibroblasts and endothelial cells. It has since been shown that IP-10 mRNA is also induced by LPS, IL-1β, TNF-α, IL-12 and viruses. Additional cell types that have been shown to express IP-10 include activated T-lymphocytes, splenocytes, keratinocytes, osteoblasts, astrocytes, and smooth muscle cells. IP-10 is also expressed in psoriatic and lepromatous lesions of skin. The mouse homologue of human IP-10, Crg-2, has been cloned andshown to share approximately 67% amino acid sequence identity with human IP-10.This chemokine elicits its effects by binding to the cell surface chemokine receptor CXCR3. IP-10 has been attributed to several roles, such as chemo attraction for monocytes and T cells, promotion of T cell adhesion to endothelial cells, antitumor activity, and inhibition of bone marrow colony formation and angiogenesis.

Protein Information

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.