

## B7-2/CD86

Catalog # PVGS1522

### Product Information

---

<b>Primary Accession Species</b>	<a href="#">P42081</a> Human
<b>Sequence</b>	Leu20-Pro247
<b>Purity</b>	> 95% as analyzed by SDS-PAGE
<b>Endotoxin Level</b>	
<b>Biological Activity</b>	Immobilized B7-2/CD86, hFc, Human at 5.0 µg/ml (100 µl/well) can bind human Biotin-CD28-Fc.
<b>Expression System</b>	HEK 293
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS, 5% trehalose and mannitol. It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH <sub>2</sub> O or PBS up to 100 µg/ml.
<b>Reconstitution</b>	
<b>Storage &amp; Stability</b>	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

### Additional Information

---

<b>Gene ID</b>	942
<b>Other Names</b>	T-lymphocyte activation antigen CD86, Activation B7-2 antigen, B70, BU63, CTLA-4 counter-receptor B7.2, FUN-1, CD86, CD86, CD28LG2
<b>Target Background</b>	B7-1 and B7-2 are homologous costimulatory ligands expressed on the surface of antigen presenting cells (APCs), both are type 1 transmembrane proteins with a membrane distal IgV and a membrane proximal IgC domain. They share ~25% sequence homology and interact with the same receptors, CD28 and CTLA-4. Binding of these molecules to the T cell costimulatory receptors, CD28 and CTLA-4, is essential for the activation and regulation of T cell immunity. T cell activation requires engagement of the T cell receptor (TCR) with the peptide-MHC complex presented on the cell surface of antigen presenting cells (APCs). In addition to this antigen-specific interaction, a second interaction involving costimulatory receptors (CD28, ICOS) on T cells and their respective ligands (B7-1/B7-2, ICOS-L) on APCs is required for optimal T cell activation. B7-1 and B7-2 may also function to deliver signal into dendritic cells. While B7-1 favors binding to CTLA-4, B7-2 shows a preference

for CD28.

## Protein Information

---

<b>Name</b>	CD86
<b>Synonyms</b>	CD28LG2
<b>Function</b>	Receptor involved in the costimulatory signal essential for T-lymphocyte proliferation and interleukin-2 production, by binding CD28 or CTLA-4 (PubMed: <a href="#">12196291</a> ). May play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation (PubMed: <a href="#">7527824</a> ). Also involved in the regulation of B cells function, plays a role in regulating the level of IgG(1) produced. Upon CD40 engagement, activates NF-kappa-B signaling pathway via phospholipase C and protein kinase C activation (By similarity).
<b>Cellular Location</b>	Cell membrane; Single-pass type I membrane protein
<b>Tissue Location</b>	Expressed by activated B-lymphocytes and monocytes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.