

# PD-1

Catalog # PVGS1554

## Product Information

---

|                                  |  |
|----------------------------------|--|
| <b>Primary Accession Species</b> | <a href="#">Q02242</a><br>Mouse  |
| <b>Sequence</b>                  | Leu25-Gln167   |
| <b>Purity</b>                    | > 95% as analyzed by SDS-PAGE  |
| <b>Endotoxin Level</b>           |  |
| <b>Expression System</b>         | HEK 293  |
| <b>Formulation</b>               | Lyophilized from a 0.2 $\mu$ m filtered solution in PBS.   |
| <b>Reconstitution</b>            | It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH <sub>2</sub> O or PBS up to 100 $\mu$ g/ml.   |
| <b>Storage &amp; Stability</b>   | Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles. |

## Additional Information

---

|                          |  |
|--------------------------|--|
| <b>Gene ID</b>           | 18566  |
| <b>Other Names</b>       | Programmed cell death protein 1, Protein PD-1, mPD-1, CD279, Pdcd1 {ECO:0000312   MGI:MGI:104879}  |
| <b>Target Background</b> | Programmed cell death protein 1, also known as PD-1 and CD279 (cluster of differentiation 279) or PDCD1, is a protein that in humans is encoded by the PDCD1 gene. PD-1 is a cell surface receptor that belongs to the immunoglobulin superfamily and is expressed on T cells and pro-B cells. PD-1 binds two ligands, PD-L1 and PD-L2. PD-1 and its ligands play an important role in down regulating the immune system by preventing the activation of T-cells, which in turn reduces autoimmunity and promotes self-tolerance. The inhibitory effect of PD-1 is accomplished through a dual mechanism of promoting apoptosis (programmed cell death) in antigen specific T-cells in lymph nodes while simultaneously reducing apoptosis in regulatory T cells (suppressor T cells). |

## Protein Information

---

|                          |  |
|--------------------------|--|
| <b>Name</b>              | Pdcd1 {ECO:0000312 MGI:MGI:104879}   |
| <b>Function</b>          | Inhibitory receptor on antigen activated T-cells that plays a critical role in induction and maintenance of immune tolerance to self (PubMed: <a href="#">10485649</a> , PubMed: <a href="#">11209085</a> , PubMed: <a href="#">11698646</a> , PubMed: <a href="#">21300912</a> ). Delivers inhibitory signals upon binding to ligands, such as CD274/PDCD1L1 and CD273/PDCD1LG2 (PubMed: <a href="#">11015443</a> , PubMed: <a href="#">11224527</a> , PubMed: <a href="#">18287011</a> , PubMed: <a href="#">18641123</a> , PubMed: <a href="#">22641383</a> ). Following T-cell receptor (TCR) engagement, PDCD1 associates with CD3-TCR in the immunological synapse and directly inhibits T-cell activation (PubMed: <a href="#">22641383</a> ). Suppresses T-cell activation through the recruitment of PTPN11/SHP-2: following ligand-binding, PDCD1 is phosphorylated within the ITSM motif, leading to the recruitment of the protein tyrosine phosphatase PTPN11/SHP-2 that mediates dephosphorylation of key TCR proximal signaling molecules, such as ZAP70, PRKCQ/PKCtheta and CD247/CD3zeta (PubMed: <a href="#">11698646</a> , PubMed: <a href="#">22641383</a> ). The PDCD1-mediated inhibitory pathway is exploited by tumors to attenuate anti-tumor immunity and facilitate tumor survival (By similarity). |
| <b>Cellular Location</b> | Cell membrane; Single-pass type I membrane protein   |
| <b>Tissue Location</b>   | Thymus-specific..  |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.