

FGL1

Catalog # PVGS1594

Product Information

Primary Accession Species	Q08830 Human
Sequence	Leu23-Ile312
Purity	> 95% as analyzed by SDS-PAGE
Endotoxin Level Biological Activity	Immobilized h_LAG3 at 5.0 μ g/ml (100 μ l/well) can bind FGL1, hFc, Human with a linear range of 0.097–50 μ g/ml when detected by Mouse Anti-Human IgG Fc-HRP.
Expression System	HEK 293
Formulation Reconstitution	Lyophilized from a 0.2 μ m filtered solution in PBS. It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μ g/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	2267
Other Names	Fibrinogen-like protein 1, HP-041, Hepassocin, HPS, Hepatocyte-derived fibrinogen-related protein 1, HFREP-1, Liver fibrinogen-related protein 1 {ECO:0000303 Ref.3}, LFIRE-1 {ECO:0000303 Ref.3}, FGL1 {ECO:0000303 PubMed:18039467, ECO:0000312 HGNC:HGNC:3695}
Target Background	FGL1 (Fibrinogen-like protein 1), also known as hepatocyte-derived fibrinogen-related protein 1 (HFREP-1), HP-041, hepassocin (HPS), and liver fibrinogen-related protein 1 (LFIRE-1), is a liver-specific secreted protein belonging to the fibrinogen superfamily whose members share a fibrinogen domain at their C-termini. FGL1 is an immune suppressive molecule that inhibits the activation of antigen-specific T cells by acting as a major ligand of LAG3, and binds LAG3 independently of MHC class II.

Protein Information

Name	FGL1 {ECO:0000303 PubMed:18039467, ECO:0000312 HGNC:HGNC:3695}
Function	Immune suppressive molecule that inhibits antigen-specific T- cell activation by acting as a major ligand of LAG3 (PubMed: 30580966). Responsible for LAG3 T-cell inhibitory function (PubMed: 30580966). Binds LAG3 independently from MHC class II (MHC-II) (PubMed: 30580966). Secreted by, and promotes growth of, hepatocytes (PubMed: 11470158 , PubMed: 19880967).
Cellular Location	Secreted. Note=Secreted in the blood plasma
Tissue Location	Under normal conditions, liver-specific.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.