

CD24

Catalog # PVGS1622

Product Information

Primary Accession Species	P24807 Mouse
Sequence	Ser27-Gly53
Purity	> 90% as analyzed by SDS-PAGE
Endotoxin Level	
Expression System	CHO 3E7
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS.
Reconstitution	It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH ₂ O or PBS up to 100 μ g/ml.
Storage & Stability	Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Additional Information

Gene ID	12484
Other Names	Signal transducer CD24, Lymphocyte antigen 52, Ly-52, M1/69-J11D heat stable antigen, HSA, Nectadrin, R13-Ag, X62 heat stable antigen, CD24, Cd24, Cd24a, Ly-52
Target Background	Signal transducer CD24 also known as cluster of differentiation 24 or heat stable antigen CD24 (HSA) is a protein that in mouse is encoded by the CD24 gene. CD24 is a cell adhesion molecule. CD24 is a sialoglycoprotein expressed at the surface of most B lymphocytes and differentiating neuroblasts. It is also expressed on neutrophils and neutrophil precursors from the myelocyte stage onwards. The encoded protein is anchored via a glycosyl phosphatidylinositol (GPI) link to the cell surface. The protein also contributes to a wide range of downstream signaling networks and is crucial for neural development. Cross-linking of CD24 on the surface of neutrophils induces apoptosis, and this appears to be defective in sepsis.

Protein Information

Name	Cd24
Synonyms	Cd24a, Ly-52
Function	May have a pivotal role in cell differentiation of different cell types. May have a specific role in early thymocyte development. Signaling could be triggered by the binding of a lectin-like ligand to the CD24 carbohydrates, and transduced by the release of second messengers derived from the GPI-anchor. Modulates B-cell activation responses (By similarity). In association with SIGLEC10 may be involved in the selective suppression of the immune response to danger- associated molecular patterns (DAMPs) such as HMGB1, HSP70 and HSP90 (PubMed: 19264983). Plays a role in the control of autoimmunity (PubMed: 20200274).
Cellular Location	Cell membrane; Lipid-anchor, GPI-anchor.
Tissue Location	In lymphoid, myeloid, and erythroid cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.